arXiv:0803.0425v1 [math.NT] 4 Mar 2008

PAIR CORRELATION OF THE ZEROS OF THE DERIVATIVE OF THE RIEMANN ξ-FUNCTION

DAVID W. FARMER AND STEVEN M. GONEK

Abstract. The complex zeros of the Riemannn zeta-function are identical to the zeros of the Riemann xi-function, $\xi(s)$. Thus, if the Riemann Hypothesis is true for the zeta-function, it is true for $\xi(s)$. Since $\xi(s)$ is entire, the zeros of $\xi'(s)$, its derivative, would then also satisfy a Riemann Hypothesis. We investigate the pair correlation function of the zeros of $\xi'(s)$ under the assumption that the Riemann Hypothesis is true. We then deduce consequences about the size of gaps between these zeros and the proportion of these zeros that are simple.

1. Introduction

Riemann's xi-function is defined by

$$\xi(s) = \frac{1}{2}s(s-1)\pi^{-s/2}\Gamma(s/2)\zeta(s), \tag{1.1}$$

where $\Gamma(s)$ is the Euler Γ-function and $\zeta(s)$ is the Riemann ζ-function. The ξ-function is an entire function of order 1 with functional equation

$$\xi(1-s) = \xi(s), \tag{1.2}$$

and its only zeros are the complex zeros of $\zeta(s)$. Thus, if the Riemann Hypothesis (RH) is true, all the zeros of $\xi(s)$ have real part one half, and the same would hold for all the zeros of the derivative $\xi'(s)$. We assume RH throughout this paper and investigate the distribution of the zeros $\rho = \frac{1}{2} + \mathrm{i}\gamma$ of $\xi'(s)$.

The distribution of zeros of ξ' is of interest for number-theoretic reasons connected to the problem of Landau-Siegel zeros, and also in connection to the general behavior of zeros of entire functions under differentiation. We discuss those motivations in the next section.

We calculate

$$F_1(\alpha, T) = N_1(T)^{-1} \sum_{0<\gamma,\gamma'\le T} T^{\mathrm{i}\alpha(\gamma-\gamma)} w(\gamma-\gamma') \ , \tag{1.3}$$

where the sum is over pairs of ordinates of zeros of $\xi'(s)$ and $w(u) = 4/(4+u^2)$ is a weight function. The normalizing factor $N_1(T) \sim \frac{1}{2\pi}T\log T$ in front of the sum is the number of zeros of ξ' with ordinates in $[0,T]$, which (on RH) differs from the number of zeros of ξ by at most 1.

Both authors were supported by the American Institute of Mathematics and the NSF Focused Research Group grant DMS 0244660. Work of the second author was also supported by NSF grants DMS 0201457 and DMS 0653809.

Theorem 1.1. *Let K be an arbitrary large positive integer. Assuming RH we have*

$$\begin{aligned} F_1(\alpha, T) &= N(T)^{-1} \sum_{0<\gamma,\gamma'\le T} T^{\mathrm{i}\alpha(\gamma-\gamma')} w(\gamma-\gamma') \\ &= (1+o(1))T^{-2|\alpha|}\log T + |\alpha| - 4|\alpha|^2 + \sum_{k=1}^{K} \frac{(k-1)!}{(2k)!}(2|\alpha|)^{2k+1} + o_K(1) \end{aligned}$$

as $T \to \infty$, for $|\alpha| < 1$.

This theorem is an analogue of Hugh Montgomery's result [16] on the pair correlation of zeros of the ζ-function. He considered the function

$$F(\alpha, T) = N(T)^{-1} \sum_{0<\gamma_0,\gamma_0'\le T} T^{\mathrm{i}\alpha(\gamma_0-\gamma_0)} w(\gamma_0-\gamma_0') \ , \tag{1.4}$$

where the sum is over pairs of ordinates of zeros of $\xi(s)$ and $w(u) = 4/(4+u^2)$. (We use γ_0 for zeros of ξ because γ refers to zeros of ξ' in this paper). In the terminology of Random Matrix Theory (RMT), $F(\alpha; T)$ is called the "2-point form factor", although sometimes it is mistakenly referred to as the pair correlation function. In fact, $F(\alpha; T)$ is the Fourier transform of the pair correlation function. Montgomery proved that $F(\alpha; T)$ has main term $T^{-2|\alpha|}\log T + |\alpha|$ for $|\alpha| < 1$. That is, $F(\alpha; T)$ behaves like a Dirac δ-function at 0 and is asymptotically $|\alpha|$ when $\varepsilon < |\alpha| < 1$. We see the same δ-like behavior in $F_1(\alpha; T)$; this is not surprising since the spike at $\alpha = 0$ is simply a consequence of the density of zeros and the weight function. The behavior of F and F_1 for $0 < |\alpha| < 1$, however, is quite different, as illustrated in Figure 1.



FIGURE 1. $F_1(\alpha; T)$ for $0 < \alpha < 0.64$ and $0 < \alpha < 1$. For comparison, the dotted line in both plots is the form factor $F(\alpha; T)$ for the zeros of the ξ-function.

Montgomery used his result on $F(\alpha; T)$ to obtain various estimates on the spacing and multiplicity of zeros of the ξ-function. Applying the same methods to Theorem 1.1 we obtain

Corollary 1.2. *Assume RH. A positive proportion of gaps between zeros of ξ' are less than 0.91 times the average spacing, and more than 3.5% of the normalized neighbor gaps between zeros of ξ' are smaller than average.*

Corollary 1.3. *Assume RH. More than 85.84% of the zeros of ξ' are simple.*

It is not surprising that the first corollary is weaker than the corresponding result for the ξ-function and the second is stronger. The reasons are discussed in the next section.

Conrey [2] has shown unconditionally that at least 79.874% of the zeros of ξ' are simple and on the critical line. Thus, Corollary 1.3 gives a conditional improvement of his result. Inserting this estimate into formula (6) of Farmer [7] improves (again on RH) the unconditional estimate there of $(0.63952 + o(1))N(T)$ for the number of distinct zeros of the ξ-function.

Corollary 1.4. *Assume RH. The number of distinct zeros of the zeta-function in* $[0, T]$ *is larger than* $(0.6544 + o(1))N(T)$.

In the next section we discuss the motivations for our work arising from the distribution of zeros of entire functions and the problem of Landau-Siegel zeros. In section 3 we state an explicit formula relating the zeros of ξ' to prime numbers, and in section 4 we begin the proof of Theorem 1.1 and identify the main terms. In section 5 we complete the proof, except for an arithmetic proposition which is proven in section 6. In section 7 we prove the explicit formula used in section 3. One important investigation we have not carried out here is the heuristic determination of $F_1(\alpha; T)$ when $|\alpha| \geq 1$.

2. The Alternative Hypothesis and the process of differentiation

Montgomery's study of the statistical behavior of zeros of the Riemann zeta-function was motivated by the problem of Landau-Siegel zeros. The idea is that a real zero very close to 1 of $L(s, \chi_d)$ would have a profound effect on the zeros of the Riemann zeta-function: in a certain range all the zeros would be on the critical line and would have a peculiar spacing. Set

$$\tilde{\gamma}_0 = \frac{1}{2\pi}\gamma_0 \log\left(\frac{\gamma_0}{2\pi}\right) \tag{2.1}$$

and denote consecutive zeros of the zeta-function by $\gamma_0 \leq \gamma_0^+$, so that $\tilde{\gamma}_0{}^+ - \tilde{\gamma}_0$ is 1 on average. The existence of a Siegel zero implies that in a certain range almost all the zeros of the zeta-function satisfy $\tilde{\gamma}_0^+ - \tilde{\gamma}_0 > \frac{1}{2} - \varepsilon$. Thus, one could disprove the existence of Landau-Siegel zeros by showing that $\tilde{\gamma}_0^+ - \tilde{\gamma}_0 \leq 0.49$, say, sufficiently often. Montgomery's result, however, only allows one to conclude that $\tilde{\gamma}_0^+ - \tilde{\gamma}_0 \leq 0.63$ a positive proportion of the time.

Montgomery refers to the connection to Landau-Siegel zeros in his paper [16], and similar connections are mentioned in unpublished work of Heath-Brown. At present the only published account is due to Conrey and Iwaniec [4]. They show that the existence of Landau-Siegel zeros implies that, in a certain range, most of the spacings between consecutive zeros of the zeta function are close to multiples of half the average spacing. That is, $\tilde{\gamma}_0^+ - \tilde{\gamma}_0$ is close to $\frac{1}{2}$ or 1, or $\frac{3}{2}$, etc. Although Conrey and Iwaniec give explicit estimates only in the case of small spacings between zeros, their main result can be used to show, for example, that if $\tilde{\gamma}_0^+ - \tilde{\gamma}_0$ was often close to 0.8, then there are no Landau-Siegel zeros. The estimates in such cases, however, have not been worked out yet.

If the statistics of the zeros of the zeta-function are governed by random matrix theory (RMT), then one could immediately conclude there are no Landau-Siegel zeros because the neighbor spacing is supported on all of $(0, \infty)$. Since there are no immediate prospects of proving that the zeros of the zeta-function follow random matrix statistics (or disproving Landau-Siegel zeros by another method), it is interesting to probe the boundary of what distributions are possible for zeros of the zeta-function. The following has been proposed as a test case:

The Alternative Hypothesis (AH) *There exists a real number* T_0 *such that if* $\gamma_0 > T_0$, *then*

$$\tilde{\gamma}_0^+ - \tilde{\gamma}_0 \in \frac{1}{2}\mathbb{Z}. \tag{2.2}$$

That is, almost all the normalized neighbor spacings are an integer or half-integer.

One can also formulate weaker versions, where the normalized spacings are approximately integers or half-integers.

2.1. **Consequences of AH.**

AH is obviously absurd, but it has not been disproven. A sufficiently strong disproof would show that there are no Landau-Siegel zeros. AH implies that Montgomery's function $F(\alpha; T)$ is periodic with period two. Thus, on AH the graph of $F(\alpha; T)$ would look like this:



Figure 2. $F(\alpha; T)$ on the Alternative Hypothesis (the heavy vertical lines represent the Dirac δ-functions at the even integers).

Similarly, the triple correlation function form factor $F(\alpha, \beta; T)$ (see [12, 18, 9]) would be periodic with period two in both α and β. Unfortunately, this does not seem to completely determine $F(\alpha, \beta; T)$ because $F(\alpha, \beta; T)$ is only known on the hexagon $|\alpha| < 1$, $|\beta| < 1$, $|\alpha - \beta| < 1$.

If we let g_μ denote the proportion of $\tilde{\gamma}_0$ such that $\tilde{\gamma}_0^+ - \tilde{\gamma}_0 = \mu$, a straightforward calculation assuming AH leads to $g_0 = 0$, $g_{\frac{1}{2}} = \frac{1}{2} - \frac{2}{\pi^2} \approx 0.297$, and $0.405 \approx \frac{4}{\pi^2} \le g_1 \le \frac{1}{2}$. Thus, one could disprove AH by showing that more than 30% of the normalized neighbor gaps of ξ are less than 0.999. We note that RMT predicts that 53.39% of the neighbor gaps are smaller than average. Montgomery's result implies that more than 12.3% of the neighbor gaps are smaller than average (set $\lambda = 1 - \varepsilon$ at the bottom of page 192 of [16]). Corollary 1.2 gives information about consecutive small gaps between zeros of ξ. This is discussed further in the next subsection.

One wonders whether AH and the existing results on zero correlations determine the distribution of neighbor spacings. To specify g_1 seems to require using the triple correlation $F(\alpha, \beta; T)$ to determine how often two consecutive normalized neighbor gaps of size $\frac{1}{2}$ can occur. So this question may be equivalent to the question of whether AH determines all the correlation functions.

Goldston and Montgomery [11] showed that the pair correlation conjecture is equivalent to

$$\int_1^X (\psi(x+h) - \psi(x) - h)^2 \, dx \sim hX \log\left(\frac{X}{h}\right), \tag{2.3}$$

for h in a certain range depending on X. Here $\psi(x) = \sum_{n\le x} \Lambda(n)$, where Λ is the von Mangoldt function: $\Lambda(n) = \log p$ if $n = p^m$ with p prime, and 0 otherwise. Montgomery and Soundararajan [17] interpret (2.3) as saying $\psi(x + h) - \psi(x)$ has mean h and variance $h \log\left(\frac{X}{h}\right)$, and they note that the Cramér model of the primes predicts a larger variance of $h \log X$. It would be interesting to see what the right-hand side of (2.3) equals if one assumes the Alternative Hypothesis.

These connections indicate the value of studying the statistics of the zeros of the zeta-function. We now explain the connection to the zeros of ξ'.

2.2. **Zeros of derivatives.** The statistics of the zeros of ξ' are interesting because of their connection with the Alternative Hypothesis and also as an illustration of the general behavior of the zeros of derivatives of an entire function upon repeated differentiation.

One motivation for studying the analogue of Montgomery's function for the zeros of ξ' is the expectation that our Theorem 1.1 might contradict AH. The zeros of ξ' are influenced by the zeros of ξ in complicated ways, so it seems unlikely that RMT and AH would predict the same behavior for $F_1(\alpha; T)$ for $|\alpha| < 1$. But there are several caveats. First, as described in the previous section, it is not known whether or not AH determines all the correlation functions of the zeros. This may lead to some flexibility in $F_1(\alpha; T)$ for $|\alpha| < 1$, which may be consistent with Theorem 1.1. Second, it is not known how to transfer a measure on the zeros of ξ to a measure on the zeros of ξ'. Thus, even if AH determined all the correlations of the zeros of ξ, it is still an unsolved problem to determine the correlations of the zeros of ξ'. Third, merely contradicting AH is not sufficient to disprove the existence of Landau-Siegel zeros. AH is an extreme example of a possible consequence of Landau-Siegel zeros. Presumably an extension of the work of Conrey and Iwaniec [4] would show that Landau-Siegel zeros imply that $F(\alpha; T)$ approximately follows Figure 2 for some range of α.

Another motivation is to understand the general behavior of zeros under differentiation. The Riemann Ξ-function is defined as $\Xi(z) = \xi(\frac{1}{2} + iz)$. The Ξ-function is an entire function of order 1 that is real on the real axis. For such functions, repeated differentiation causes the zeros to migrate to the real axis [6, 15, 14]. Thus, in any bounded region the Riemann Hypothesis is true for the nth derivative $\Xi^{(n)}(z)$ for sufficiently large n.

It is conjectured [8] that for real entire functions of order 1, whose zeros lie in a strip around the real axis, not only do the zeros migrate toward the real axis, but they also approach equal spacing. That is, the derivatives approach a multiple of $e^{ax} \cos(bx + c)$. This conjecture has been proven with some restrictions on the distribution of zeros [8] and for some special cases, such as the Ξ-function [13] and the reciprocal of the gamma function [1].

The reason differentiation leads to equally spaced zeros is that, locally, the zeros of f' move away from concentrations of zeros of f and towards regions with fewer zeros of f. Thus, small gaps become larger and large gaps become smaller. Figure 3 illustrates these ideas.

A consequence is that if ξ' has a small gap between consecutive zeros, then ξ must have had an even smaller gap. Thus, one should expect that the existence of Landau-Siegel zeros could be disproven by showing that ξ' has sufficiently many zeros separated by less than half the average spacing. That is, one should be able to extend the result of Conrey-Iwaniec [4] to the zeros of $\xi^{(n)}$ for any $n \ge 1$. However, we have not worked out the necessary details to show that the required number of small gaps between zeros of ξ' leads to the same number of small gaps between the zeros of ξ. Thus, at present we just mention this as a motivating principle. See [8] for a discussion.



FIGURE 3. The circles are the midpoints of neighboring zeros of f, and the dots are the zeros of f'.

Based on these ideas, we see that Corollary 1.2 does not contradict the Alternative Hypothesis. On average each normalized gap of size $\frac{1}{2}$ is adjacent to a gap of size $\frac{1}{2}$ or 1, so AH implies that at least 29.7% of the normalized neighbor gaps of ξ' are smaller than 1. If Corollary 1.2 could be improved to show that ξ' has normalized zero gaps smaller than 0.75, then that would imply, on AH, that ξ has consecutive gaps of size $\frac{1}{2}$.

Our final motivation is to understand the manner in which differentiation causes the zeros to become equally spaced. Let F_n denote the analogue of F_1 involving the zeros of $\xi^{(n)}(z)$ rather than the zeros of $\xi'(z)$. Since the zeros of $\xi^{(n)}(z)$ approach equal spacing as n increases, F_n approaches a sum of Dirac δ-functions supported at the integers. We would like to understand this transition to a sum of δ-functions. The only case we know of where this has been worked out explicitly is for random trigonometric polynomials [10]. Theorem 1.1 is a first step in this direction, and Figure 1 does seem to illustrate the expected behavior. In a recent Ph.D. thesis, Jim Bian has now worked out explicit formulas for $F_n(\alpha; T)$ for $n \geq 2$ and $|\alpha| < 1$.

2.3. **Connection to RMT.**

It would be interesting to know the random matrix analogue of Theorem 1.1. At present, this is one of the few calculations carried out for the Riemann zeta-function for which a random matrix analogue seems out of reach. One issue is that in the random matrix world there seems to be no direct analogue of the Riemann ξ-function. See Section 1.2 of [3] for a discussion. The closest match is

$$z^{-n/2}\Lambda(z), \tag{2.4}$$

where $\Lambda(z)$ is the characteristic polynomial of a random matrix from the unitary group $U(n)$, chosen uniformly with respect to Haar measure. However, this is more nearly an analogue of the Hardy Z-function defined by $Z(t) = \chi(\frac{1}{2}+it)^{-\frac{1}{2}}\zeta(\frac{1}{2}+it)$. This was the motivation of Conrey and Soundararajan [5], who did similar calculations to ours for the Z-function. But it turns out that the zeros of $\Xi'(t)$ and $Z'(t)$ should have similar correlation functions. Since $\Xi(t)$ is approximately $e^{-\frac{\pi t}{2}}Z(t)$, the corresponding zeros of $\Xi'(t)$ and $Z'(t)$ generally differ by $O(1/\log^2 t)$. So one should expect the correlation functions of their zeros to be equal, to leading order. We provide a more rigorous explanation in Section 7.2. And those correlation functions should equal, to leading order, the correlation function of the zeros of the derivative of the completed characteristic polynomial (2.4).

3. An explicit formula and first steps of the proof

The general outline of our calculation is similar to the case of zeros of the zeta-function. We start with an explicit formula relating a sum over zeros of ξ' to a sum involving number-theoretic expressions, and then compute the mean-square of both sides. Our main complication is that the number-theoretic side of the explicit formula is not a Dirichlet series, but an "approximate" Dirichlet series, by which we mean a Dirichlet series whose coefficients may depend on s.

The explicit formula is derived in Section 7. Here we just outline the calculation and state the formula.

We begin by noting that $\xi''/\xi'(s)$ has simple poles with residue 1 at the zeros of $\xi'(s)$ and no others; here zeros are counted as many times as their multiplicity. Our first goal is to write $\xi''/\xi'(s)$ as an "approximate" Dirichlet series. From the definition of the ξ-function (1.1) we have

$$\frac{\xi'}{\xi}(s) = L(s) + \frac{\zeta'}{\zeta}(s)\ , \tag{3.1}$$

where

$$\begin{aligned} L(s) &= \frac{1}{s} + \frac{1}{s-1} - \frac{\log \pi}{2} + \frac{1}{2}\frac{\Gamma'}{\Gamma}\left(\frac{s}{2}\right) \\ &= \frac{1}{2}\log\frac{s}{2\pi} + O\left(\frac{1}{|s|+2}\right) \end{aligned} \tag{3.2}$$

and

$$L'(s) \ll \frac{1}{|s|+2} \tag{3.3}$$

in the region $|\arg s| < \pi - \delta$, $|s| \geq 1/4$, say.

Multiplying both sides of (3.1) by $\xi(s)$ and calculating the logarithmic derivative leads to the following lemma, the proof of which is in Section 7.

Lemma 3.1. *For $\sigma \geq 1+\varepsilon$, $|t| \geq T_\varepsilon$, and K a large positive integer we have*

$$\frac{\xi''}{\xi'}(s) = L(s) + \sum_{n=1}^{\infty} \frac{a_K(n,s)}{n^s} + O\left(\frac{1}{\varepsilon^2 2^K}\right)\ . \tag{3.4}$$

Here we have written

$$a_K(n,s) = \sum_{k=0}^{K} \frac{\alpha_k(n)}{L(s)^k}, \tag{3.5}$$

where

$$\alpha_j(n) = \begin{cases} -\Lambda(n) & \text{if} \quad j = 0, \\ \Lambda_{j-1} * \Lambda\log(n) & \text{if} \quad j \geq 1. \end{cases} \tag{3.6}$$

The function Λ_j for $j \geq 0$ is the j-fold convolution of the von Mangoldt function, defined by

$$\left(-\frac{\zeta'}{\zeta}(s)\right)^j = \sum_{n=1}^{\infty} \frac{\Lambda_j(n)}{n^s}\,. \tag{3.7}$$

for $\sigma > 1$. Simple estimates of Λ_j and α_k will be used repeatedly. We have the trivial bound

$$\Lambda_j(n) \leq (\log n)^j \qquad (j = 1, 2, \dots)\,, \tag{3.8}$$

which follows from iterating

$$\Lambda_j(n) = \sum_{d|n} \Lambda(d)\Lambda_{j-1}\left(\frac{n}{d}\right) \leq \left(\max_{d|n} \Lambda_{j-1}\left(\frac{n}{d}\right)\right) \sum_{d|n} \Lambda(d) \leq \Lambda_{j-1}(n^*) \log n \tag{3.9}$$

for some divisor n^* of n. Similarly,

$$\alpha_k(n) = \Lambda_{k-1} * \Lambda \log(n) \leq (\log n)^{k+1} \qquad (k = 1, 2, \dots)\,, \tag{3.10}$$

from which it follows that

$$|a_K(n, s)| \leq \log n \sum_{k=0}^{K} \left(\frac{\log n}{|L(s)|}\right)^k \ll \log^{K+1} n \tag{3.11}$$

when $-1 \leq \sigma \leq 2$, say.

A contour integral of ξ''/ξ' times an appropriate kernel leads to the following explicit formula; the details are in Section 7.

Proposition 3.2. *For $5/4 < \sigma < 2$, $x \geq 1$, $0 < \varepsilon < 1/8$, and K a positive integer*

$$\begin{aligned} &(2\sigma - 1) \sum_{\gamma} \frac{x^{\mathrm{i}\gamma}}{(\sigma - 1/2)^2 + (t - \gamma)^2} \\ &= x^{-1/2} \left(\sum_{n \leq x} a_K(n, 1 - \bar{s}) \left(\frac{x}{n}\right)^{1-\bar{s}} + \sum_{n > x} a_K(n, s) \left(\frac{x}{n}\right)^s \right) \\ &\quad + x^{1/2-\bar{s}} \log \frac{\tau}{2\pi} + O(x^{1/2-\sigma}) + O_{\varepsilon,K}\left(x^{1/2}\tau^{-1} \max(x^{\varepsilon}, \log^{2K+2} x)\right)\,, \end{aligned} \tag{3.12}$$

where $\tau = |t| + 2$.

4. BEGINNING OF THE PROOF

Set $\sigma = 3/2$ in Proposition 3.2, write the resulting equation as $L(x, t) = R_1(x, t) + R_2(x, t) + R_3(x, t) + R_4(x, t)$, and calculate

$$\int_0^T |L(x, t)|^2 \mathrm{d}t = \int_0^T |R_1(x, t) + R_2(x, t) + R_3(x, t) + R_4(x, t)|^2 \mathrm{d}t\,. \tag{4.1}$$

The left-hand side may be treated in exactly the same way as the corresponding expression in Montgomery [16], to whom we refer the reader (cf. pp. 187–188). We find

$$\int_0^T |L(x, t)|^2 \mathrm{d}t = 2\pi \sum_{0<\gamma,\gamma' \leq T} x^{\mathrm{i}(\gamma-\gamma')} w(\gamma - \gamma') + O(\log^3 T)\,, \tag{4.2}$$

where $w(u) = 4/(4 + u^2)$.

Next we begin the calculation of the right-hand side of (4.1). In various ranges of x, one or another of the integrals $\int_0^T |R_i(x,t)|^2 \mathrm{d}t$ dominates the others, and we record the following useful formula for later. Given x, let $\mathcal{R}_{\mathbf{1}}(x)$ be the largest of $\int_0^T |R_i(x,t)|^2 \mathrm{d}t$ ($i = 1,2,3,4$), $\mathcal{R}_{\mathbf{2}}(x)$ the next largest, and so on. Then we have

$$\int_0^T |R_1(x,t) + R_2(x,t) + R_3(x,t) + R_4(x,t)|^2 \mathrm{d}t = \mathcal{R}_{\mathbf{1}}(x) + O\left((\mathcal{R}_{\mathbf{1}}(x)\mathcal{R}_{\mathbf{2}}(x))^{1/2} \right) \tag{4.3}$$

We find

$$\int_0^T |R_2(x,t)|^2 \mathrm{d}t = \int_0^T \left| x^{-1+\mathrm{i}t} \log \frac{\tau}{2\pi} \right|^2 \mathrm{d}t = \frac{T}{x^2}\left(\log^2 T + O(\log T) \right) , \tag{4.4}$$

$$\int_0^T |R_3(x,t)|^2 \mathrm{d}t \ll \frac{T}{x^2} \tag{4.5}$$

and

$$\begin{aligned} \int_0^T |R_4(x,t)|^2 \mathrm{d}t &\ll_{\varepsilon,K} x \max(x^{2\varepsilon}, (\log x)^{4K+4}) \int_0^T \frac{\mathrm{d}t}{\tau^2} \\ &\ll_{\varepsilon,K} x \max(x^{2\varepsilon}, (\log x)^{4K+4}) . \end{aligned} \tag{4.6}$$

To estimate $\int_0^T |R_1(x,t)|^2 \mathrm{d}t$, we need the following

Lemma 4.1. *Let $\sigma = -1/2$ or $3/2$. Then for $x > 0$ we have*

$$\int_0^T \frac{x^{\mathrm{i}t}}{\overline{L(s)}^{\,k} L(s)^{\ell}} \, \mathrm{d}t = \begin{cases} T\left(\frac12 \log\left(\frac{T}{2\pi}\right)\right)^{-(k+l)} \left(1 + O_K\left(\frac{1}{\log T}\right)\right) & \text{if } x = 1 , \\ O_K\left(\frac{1}{|\log x|}\right) & \text{if } x \neq 1 , \end{cases} \tag{4.7}$$

where $k, \ell = 0,1,2,\ldots,K$. The same result holds for

$$\int_0^T \frac{x^{\mathrm{i}t}}{\overline{L(1-\bar{s})}^{\,k} L(s)^{\ell}} \, \mathrm{d}t . \tag{4.8}$$

Proof. We prove (4.7) only, as the proof of (4.8) is almost identical.

First observe that from (3.2) we have

$$\frac{1}{\overline{L(s)}^{\,k} L(s)^{\ell}} = \frac{1}{\left(\frac12 \log \tau/2\pi\right)^{k+l}} \left(1 + O_K\left(\frac{1}{\log \tau} \right) \right) . \tag{4.9}$$

The case $x = 1$ follows immediately.

Now suppose that $x > 0$, but $x \neq 1$. Integrating by parts, we find that our integral equals

$$\left. \frac{x^{\mathrm{i}t}}{\mathrm{i}\log x \, \overline{L(s)}^{\,k} L(s)^{\ell}} \right|_0^T + \frac{1}{\mathrm{i}\log x} \int_0^T \frac{x^{\mathrm{i}t}}{\overline{L(s)}^{\,k} L(s)^{\ell}} \left(k \frac{\overline{L'}(s)}{\overline{L}(s)} + \ell \frac{L'(s)}{L(s)} \right) \mathrm{d}t .$$

By (3.2), (3.3), and (4.9), this is

$$\begin{aligned} &\ll \frac{1}{|\log x|} \left(1 + (k+\ell) \int_0^T \frac{1}{\tau (1/2 \log \tau/2\pi)^{k+\ell+1}} \mathrm{d}t \right) \\ &\ll_K \frac{1}{|\log x|} . \end{aligned}$$

This completes the proof.

$\square$

We now come to the term

$$\begin{aligned}\int_0^T |R_1(x,t)|^2 \mathrm{d}t &= \frac{1}{x}\int_0^T \left|\sum_{n\le x} a_K(n,-1/2+\mathrm{i}t)\left(\frac{x}{n}\right)^{-1/2+\mathrm{i}t} + \sum_{n>x} a_K(n,3/2+\mathrm{i}t)\left(\frac{x}{n}\right)^{3/2+\mathrm{i}t}\right|^2 \mathrm{d}t\\
&= \frac{1}{x^2}\int_0^T \left|\sum_{n\le x} a_K(n,-1/2+\mathrm{i}t) n^{1/2-\mathrm{i}t}\right|^2 \mathrm{d}t + x^2\int_0^T \left|\sum_{n>x} a_K(n,3/2+\mathrm{i}t) n^{-3/2-\mathrm{i}t}\right|^2 \mathrm{d}t\\
&\quad + 2Re\int_0^T \overline{\left(\sum_{n\le x} a_K(n,-1/2+\mathrm{i}t) n^{1/2-\mathrm{i}t}\right)}\left(\sum_{m>x} a_K(m,3/2+\mathrm{i}t) m^{-3/2+\mathrm{i}t}\right)\mathrm{d}t\\
&= \frac{1}{x^2}R_{1,1} + x^2 R_{1,2} + 2\,\mathrm{Re}\,R_{1,3}\ ,\end{aligned}$$

say. Recalling that $a_K(n,s) = \sum_{k=0}^K \alpha_k(n)/L(s)^k$, we see by Lemma 4.1 that

$$\begin{aligned}R_{1,1} &= \sum_{m,n\le x}\sqrt{mn}\sum_{k,\ell=0}^K \overline{\alpha_k(n)}\alpha_\ell(m)\int_0^T \frac{(n/m)^{it}}{\overline{L(-1/2+it)^k}L(-1/2+it)^\ell}\,dt\\
&= (1+o(1))T\sum_{n\le x} n\left(\sum_{k,\ell=0}^K \frac{\overline{\alpha_k(n)}\alpha_\ell(n)}{(1/2\log T/2\pi)^{k+\ell}}\right)\\
&\quad + O_K\left(\sum_{1\le m<n\le x}\sqrt{mn}\left(\sum_{k,\ell=0}^K \frac{|\alpha_k(n)\alpha_\ell(m)|}{\log n/m}\right)\right)\\
&= (1+o(1))T\sum_{n\le x} n\left|\sum_{k=0}^K \frac{\alpha_k(n)}{(1/2\log T/2\pi)^k}\right|^2\\
&\quad + O_K\left(\sum_{1\le m<n\le x}\frac{\sqrt{mn}}{\log n/m}\left(\sum_{k=0}^K |\alpha_k(n)|\right)\left(\sum_{k=0}^K |\alpha_k(m)|\right)\right)\ .\end{aligned}$$

From the standard inequality

$$\sum_{1\le m<n\le x}\frac{|b(m)b(n)|}{(mn)^\sigma \log n/m} \ll x\log x\sum_{n\le x}\frac{|b(n)|^2}{n^{2\sigma}} \tag{4.10}$$

and (3.10) we see that the O-term is

$$\begin{aligned}&\ll_K x\log x\sum_{n\le x} n\left(\sum_{k=0}^K |\alpha_k(n)|\right)^2\\
&\ll_K x\log x\sum_{n\le x} n(\log n)^{2K+2}\\
&\ll_K x^3(\log x)^{2K+3}\ .\end{aligned}$$

Thus, we have

$$R_{1,1} = (1+o(1))T\sum_{n\le x} n\left|\sum_{k=0}^{K}\frac{\alpha_k(n)}{(1/2\log T/2\pi)^k}\right|^2 + O_K\left(x^3(\log x)^{2K+3}\right) .$$

Similarly, we find that

$$R_{1,2} = (1+o(1))T\sum_{n>x} n^{-3}\left|\sum_{k=0}^{K}\frac{\alpha_k(n)}{(1/2\log T/2\pi)^k}\right|^2 + O_K\left(x^{-1}(\log x)^{2K+3}\right) .$$

For $R_{1,3}$ we have

$$\begin{aligned}
R_{1,3} &= \sum_{n\le x} n^{1/2}\sum_{m>x} m^{-3/2}\sum_{k,\ell=0}^{K}\overline{\alpha_k(n)}\alpha_\ell(m)\int_0^T \frac{(mn)^{it}}{\overline{L(-1/2+it)}^{\,k}L(3/2+it)^\ell}\,dt\\
&\ll \sum_{n\le x} n^{1/2}\sum_{m>x} m^{-3/2}(\log mn)^{-1}\left(\sum_{k,\ell=0}^{K}|\alpha_k(n)\alpha_\ell(m)|\right)\\
&\ll_K \sum_{n\le x}\sum_{m>x}\frac{n^{1/2}m^{-3/2}}{\log mn}(\log m)^{K+1}(\log n)^{K+1}\\
&\ll_K \frac{1}{\log x}\left(\sum_{n\le x} n^{1/2}(\log n)^{K+1}\right)\left(\sum_{m>x} m^{-3/2}(\log m)^{K+1}\right)\\
&\ll_K x(\log x)^{2K+1} .
\end{aligned}$$

Combining these estimates, we have

$$\begin{aligned}
\int_0^T |R_1(x,t)|^2\mathrm{d}t =&(1+o(1))Tx^{-2}\sum_{n\le x} n\left|\sum_{k=0}^{K}\frac{\alpha_k(n)}{(1/2\log T/2\pi)^k}\right|^2\\
&+(1+o(1))Tx^2\sum_{n>x}\frac{1}{n^3}\left|\sum_{k=0}^{K}\frac{\alpha_k(n)}{(1/2\log T/2\pi)^k}\right|^2\\
&+O_K(x(\log x)^{2K+3}) .
\end{aligned}\tag{4.11}$$

In the next section we complete the proof of Theorem 1.1, subject to an arithmetic proposition which we prove in Section 6.

5. Proof of Theorem 1.1

It remains to evaluate (4.11) and to put the expressions in the form of Theorem 1.1. The main terms from (4.11) can be obtained from a Stieltjes integral involving

$$A(x) = A(x,K,T) = \sum_{n\le x}\left|\sum_{k=0}^{K}\frac{\alpha_k(n)}{\mathcal{L}^k}\right|^2 .\tag{5.1}$$

Here α_k is defined in (3.6) and $\mathcal{L} = \frac{1}{2}\log\frac{T}{2\pi}$. We write

$$A(x) = \sum_{k=0}^{K} A_{k,k}(x)\mathcal{L}^{-2k} + 2\sum_{k=1}^{K}\sum_{0\le \ell<k} A_{k,\ell}(x)\mathcal{L}^{-(k+\ell)}, \tag{5.2}$$

where

$$A_{k,\ell}(x) = \sum_{n\le x}\alpha_k(n)\alpha_\ell(n). \tag{5.3}$$

The following proposition, which is proven in Section 6, is sufficient to evaluate the leading order asymptotics of (4.11).

Proposition 5.1. *We have*

$$A_{k,0}(x) = \begin{cases} x\log x + O(x) & \text{if} \quad k=0\,, \\ -x\log^2 x + O(x\log x) & \text{if} \quad k=1\,, \\ O\big(x^{\frac{1}{2}+\varepsilon}\big) & \text{if} \quad k\ge 2\,. \end{cases} \tag{5.4}$$

and

$$A_{k,1}(x) = \begin{cases} x\log^3 x + O(x\log^2 x) & \text{if} \quad k=1\,, \\ O\big(x^{\frac{1}{2}+\varepsilon}\big) & \text{if} \quad k\ge 2\,. \end{cases} \tag{5.5}$$

If $k \ge 1$, then

$$A_{k,k}(x) = 2\frac{(k-1)!}{(2k)!}x\log^{2k+1}x + O(x\log^{2k}x), \tag{5.6}$$

and if $k > \ell \ge 2$, then

$$A_{k,\ell}(x) \ll x\log^{k+\ell}x. \tag{5.7}$$

As a consequence, we have

$$A(x) = x\log x\left(1 - 2\left(\frac{\log x}{\mathcal{L}}\right) + 2\sum_{k=1}^{K}\frac{(k-1)!}{(2k)!}\left(\frac{\log x}{\mathcal{L}}\right)^{2k}\right) + O_K(x) \tag{5.8}$$

for $x = T^\alpha$ with $0 < \alpha \le C_0$, where C_0 is any fixed positive number.

Proof of Theorem 1.1. We first evaluate $\int_0^T |R_1(x,t)|^2\mathrm{d}t$ given by (4.11). We have

(5.9)

$$\int_0^T |R_1(x,t)|^2\mathrm{d}t = (1+o(1))T\left\{x^{-2}\int_{1^-}^{x} u\,dA(u) + x^2\int_x^\infty u^{-3}dA(u)\right\} + O_K\left(x(\log x)^{2K+3}\right)$$

$$= (1+o(1)T\left\{-x^{-2}\int_1^x A(u)du + 3x^2\int_x^\infty u^{-4}A(u)du\right\} + O_K\left(x(\log x)^{2K+3}\right)\ . \tag{5.10}$$

Note that the boundary terms canceled when we integrated by parts.

The typical term in the series for $A(u)$ has the form $Cu(\log u)^{m+1}/\mathcal{L}^m$ and, assuming that $m \ll K$, we have

$$\frac{1}{\mathcal{L}^m}\int_1^x u(\log u)^{m+1}du = \frac{1}{\mathcal{L}^m}\frac{x^2}{2}(\log x)^{m+1}\left(1 + O_K\left(\frac{1}{\log x}\right)\right)$$

and

$$\frac{1}{\mathcal{L}^m}\int_x^\infty u^{-3}(\log u)^{m+1}du = \frac{1}{\mathcal{L}^m}\frac{1}{2x^2}(\log x)^{m+1}\left(1+O_K\left(\frac{1}{\log x}\right)\right) \ .$$

Using the first formula and (5.8), we have

$$\begin{aligned}-x^{-2}\int_1^x A(u)du =& - x^{-2}\int_1^x u\left(\log u - 2\frac{\log^2 u}{\mathcal{L}} + 2\sum_{k=1}^K \frac{(k-1)!}{(2k)!}\frac{(\log u)^{2k+1}}{\mathcal{L}^{2k}}\right)du + O_K(1)\\ =& \log x\left\{-\frac{1}{2}+\frac{\log x}{\mathcal{L}} - \sum_{k=1}^K \frac{(k-1)!}{(2k)!}\left(\frac{\log x}{\mathcal{L}}\right)^{2k}\right\}\left(1+O_K\left(\frac{1}{\log x}\right)\right) + O_K(1)\,.\end{aligned}$$

For $x \ll T^C$, the $O_K(1/\log x)$ term contributes no more than $O_K(1)$ to this. Hence,

$$-x^{-2}\int_1^x A(u)du = \log x\left(-\frac{1}{2}+\frac{\log x}{\mathcal{L}} - \sum_{k=1}^K \frac{(k-1)!}{(2k)!}\left(\frac{\log x}{\mathcal{L}}\right)^{2k}\right) + O_K(1)\ . \tag{5.11}$$

Similarly, using the second formula and (5.8), we obtain

$$\begin{aligned}3x^2\int_x^\infty u^{-4}A(u)du =& 3x^2\int_x^\infty u^{-3}\left(\log u - \frac{2\log^2 u}{\mathcal{L}} + 2\sum_{k=1}^K \frac{(k-1)!}{(2k)!}\frac{(\log u)^{2k+1}}{\mathcal{L}^{2k}}\right)du + O_K(1)\\ =& \log x\left\{\frac{3}{2} - 3\frac{\log x}{L} + 3\sum_{k=1}^K \frac{(k-1)!}{(2k)!}\left(\frac{\log x}{L}\right)^{2k}\right\} + O_K(1)\,.\end{aligned} \tag{5.12}$$

Combining terms we obtain

$$\begin{aligned}\int_0^T |R_1(x,t)|^2\mathrm{d}t =& (1+o(1))\,T\log x\left(1-2\frac{\log x}{\mathcal{L}} + 2\sum_{k=1}^K \frac{(k-1)!}{(2k)!}\left(\frac{\log x}{\mathcal{L}}\right)^{2k}\right)\\ &+ O_K(T) + O_K\left(x(\log x)^{2K+3}\right)\end{aligned} \tag{5.13}$$

provided $x \ll T^{C_0}$.

Recall from (4.1) and (4.2) that we write

$$2\pi\sum_{0<\gamma,\gamma'\le T} x^{\mathrm{i}(\gamma-\gamma')}w(\gamma-\gamma') = \mathcal{R}_1(x) + O\left((\mathcal{R}_1(x)\mathcal{R}_2(x))^{1/2}\right) + O(\log^3 T)\,, \tag{5.14}$$

where, for a given x, $\mathcal{R}_1(x)$ is the largest of $\int_0^T |R_i(x,t)|^2\mathrm{d}t$ $(i=1,2,3,4)$ and $\mathcal{R}_2(x)$ is the next largest. Now, from the various estimates we see that our $\mathcal{R}_1(x)$ term is given by

$$\begin{aligned}(1+o(1))\frac{T}{x^2}\log^2 T \qquad & \text{if } 1\le x\le(\log T)^{3/4}\ ,\\ o(T\log T) \qquad & \text{if } \quad (\log T)^{3/4} < x \le (\log T)^{3/2}\ ,\end{aligned}$$

and by

$$(1+o_K(1))\,T\log x\left(1-2\frac{\log x}{\mathcal{L}} + 2\sum_{k=1}^K\frac{(k-1)!}{(2k)!}\left(\frac{\log x}{\mathcal{L}}\right)^{2k}\right) \qquad \text{if} \quad (\log T)^{3/2} < x \le T^{1-\varepsilon}\ .$$

In each of these ranges, it happens that the $\mathcal{R}_2(x)$ term is $o(\mathcal{R}_1(x))$. Hence, taking $x = T^\alpha$ in (5.14), we find that for $0 < \alpha < 1$ and T large,

$$\begin{aligned}&2\pi \sum_{0<\gamma,\gamma'\le T} T^{\mathrm{i}\alpha(\gamma-\gamma')} w(\gamma-\gamma')\\ &= (1+o(1))T^{1-2\alpha}\log^2 T + (1+o_K(1))\alpha T\log T \left(1 - 4\alpha + 2\sum_{k=1}^{K} \frac{(k-1)!}{(2k)!}\,(2\alpha)^{2k}\right) + o_K(1)\ .\end{aligned}$$

Using (1.3) and noting that $F_1(\alpha, T)$ is an even function of α, we have proved Theorem 1.1. □

6. Proof of Proposition 5.1

We prove Proposition 5.1, which is the arithmetic portion of the calculation. We first reduce $A_{k,\ell}$ to a sum involving the arithmetic functions Λ_j. In Section 6.2 we state some lemmas which are needed in the calculation, and in Section 6.3 we evaluate the sums of the Λ_j. Then we complete the proof of Proposition 5.1 in Section 6.4

6.1. **Reduction of $A_{k,\ell}$.**

Recall that

$$A_{k,\ell}(x) = \sum_{n\le x} \alpha_k(n)\alpha_\ell(n), \tag{6.1}$$

where α_k is given by (3.6).

Lemma 6.1. *We have*

$$A_{k,0}(x) = \begin{cases} x\log x + O(x) & \text{if} \quad k=0\,, \\ -x\log^2 x + O(x\log x) & \text{if} \quad k=1\,, \\ O\big(x^{\frac12+\varepsilon}\big) & \text{if} \quad k\ge 2\,. \end{cases} \tag{6.2}$$

and

$$A_{k,1}(x) = \begin{cases} x\log^3 x + O(x\log^2 x) & \text{if} \quad k=1\ , \\ O\big(x^{\frac12+\varepsilon}\big) & \text{if} \quad k\ge 2\ . \end{cases} \tag{6.3}$$

For $2 \le \ell \le k$ we have

$$\begin{aligned} A_{k,\ell}(x) = (k-1)(\ell-1)&\sum_{p\le x}\sum_{q\le \frac{x}{p}} \log^3 p\, \log^3 q \left(\sum_{m\le \frac{x}{pq}} \Lambda_{k-2}(m)\Lambda_{\ell-2}(m)\right)\\ &+ \sum_{p\le x}\log^4 p \left(\sum_{m\le\frac{x}{p}} \Lambda_{k-1}(m)\Lambda_{\ell-1}(m)\right)\\ &+ O(x\log^{k+\ell} x). \end{aligned} \tag{6.4}$$

Proof. The form of $\alpha_k(n)$ is different when $k=0$ and 1 from what it is for larger k, and this will be reflected in our estimates for $A_{k,\ell}(x)$. We therefore treat these cases separately.

First consider the case of $A_{k,0}(x)$. By the prime number theorem we have

$$A_{0,0}(s) = \sum_{n\le x} \Lambda^2(n) = x\log x + O(x) \tag{6.5}$$

and

$$A_{1,0}(x) = -\sum_{n\le x} \Lambda^2(n)\log n = -x\log^2 x + O(x\log x)\,. \tag{6.6}$$

For $k \ge 2$ we have

$$\begin{aligned} A_{k,0}(x) &= -\sum_{n\le x} \Lambda(n)(\Lambda_{k-1} * \Lambda\log)(n) \\ &= -\sum_{p^a\le x} \log(p)(\Lambda_{k-1} * \Lambda\log)(p^a) \\ &\ll \sum_{\substack{p^a\le x\\ a\ge k}} p^{a\varepsilon} \\ &\ll x^{\frac{1}{2}+\varepsilon}. \end{aligned} \tag{6.7}$$

We have used the fact that $(\Lambda_{k-1} * \Lambda\log)(n) \ll n^{\varepsilon}$ and that this function vanishes unless n is a product of at least k (not necessarily distinct) primes and $k \ge 2$.

Next we consider $A_{k,1}(x)$ for $k \ge 1$. By the prime number theorem,

$$A_{1,1}(x) = \sum_{n\le x} (\Lambda(n)\log n)^2 = x\log^3 x + O(x\log^2 x)\ . \tag{6.8}$$

If $k \ge 2$ then

$$\begin{aligned} A_{k,1}(x) &= \sum_{n\le x} \Lambda(n)\log n\ \left(\Lambda_{k-1} * \Lambda\log\right)(n) \\ &\ll x^{\frac{1}{2}+\varepsilon}, \end{aligned} \tag{6.9}$$

exactly as in (6.7).

We now come to the general case of $A_{k,\ell}(x)$ with $k \ge \ell \ge 2$. We have

$$\begin{aligned} A_{k,\ell}(x) &= \sum_{n\le x} \alpha_k(n)\alpha_l(n) \\ &= \sum_{n\le x} \left(\sum_{p^a|n} a\log^2 p\ \Lambda_{k-1}\left(\frac{n}{p^a}\right)\right)\left(\sum_{q^b|n} b\log^2 q\ \Lambda_{\ell-1}\left(\frac{n}{q^b}\right)\right) \\ &= \sum_{n\le x} \left(\sum_{p|n} \log^2 p\ \Lambda_{k-1}\left(\frac{n}{p}\right)\right)\left(\sum_{q|n} \log^2 q\ \Lambda_{\ell-1}\left(\frac{n}{q}\right)\right) \\ &\qquad + O(x\log^{k+\ell} x) \\ &= \sum_{p\le x}\sum_{\substack{q\le \frac{x}{p}\\ q\ne p}} \log^2 p\log^2 q\left(\sum_{\substack{n\le x\\ pq|n}} \Lambda_{k-1}\left(\frac{n}{p}\right)\Lambda_{\ell-1}\left(\frac{n}{q}\right)\right) \end{aligned}$$

$$\begin{aligned}
&+\sum_{p\le x}\log^4 p\left(\sum_{\substack{n\le x\\ p|n}}\Lambda_{k-1}\left(\frac{n}{p}\right)\Lambda_{\ell-1}\left(\frac{n}{p}\right)\right)\\
&+O(x\log^{k+\ell}x)\\
&=A_1+A_2+O(x\log x^{k+\ell}x)\ , \qquad (6.10)
\end{aligned}$$

say. The estimate on the third line above was done as follows. By symmetry, it is sufficient to estimate

$$\sum_{\substack{p^a\le x\\ a\ge 1}}\sum_{\substack{q^b\le \frac{x}{p^a}\\ b\ge 2}} ab\log^2 p\ \log^2 q\left(\sum_{m\le\frac{x}{p^aq^b}}\Lambda_{k-1}(mq^b)\ \Lambda_{\ell-1}(mp^a)\right)\ . \tag{6.11}$$

By the trivial bound (3.8) this is

$$\begin{aligned}
&\ll\sum_{\substack{p^a\le x\\ a\ge 1}}\sum_{\substack{q^b\le \frac{x}{p^a}\\ b\ge 2}} ab\log^2 p\ \log^2 q\left(\sum_{m\le\frac{x}{p^aq^b}}(\log mq^b)^{k-1}\ (\log mp^a)^{\ell-1}\right)\\
&\ll x(\log x)^{k+\ell-2}\sum_{\substack{p^a\le x\\ a\ge 1}}\frac{a\log^2 p}{p^a}\sum_{\substack{q^b\le \frac{x}{p^a}\\ b\ge 2}}\frac{b\log^2 q}{q^b}\\
&\ll x(\log x)^{k+\ell-2}\sum_{\substack{p^a\le x\\ a\ge 1}}\frac{a\log^2 p}{p^a}\\
&\ll x(\log x)^{k+\ell}, \qquad (6.12)
\end{aligned}$$

as claimed. One can use Lemma 6.2 on the inner summand of (6.11) to improve this bound by a power of $\log x$, but this will not affect our final result.

It is clear that A_2 equals the second main term in (6.4), so it remains to put A_1 in the appropriate form. The point is that if p is prime and $p\nmid m$ then

$$\Lambda_k(pm)=k(\log p)\Lambda_{k-1}(m). \tag{6.13}$$

See the proof of Lemma 6.2. Thus

$$\begin{aligned}
A_1&=(k-1)(\ell-1)\sum_{p\le x}\sum_{\substack{q\le\frac{x}{p}\\ q\ne p}}\log^3 p\ \log^3 q\left(\sum_{\substack{m\le\frac{x}{pq}\\ (m,pq)=1}}\Lambda_{k-2}(m)\Lambda_{\ell-2}(m)\right)\\
&+\sum_{p\le x}\sum_{\substack{q\le\frac{x}{p}\\ q\ne p}}\log^2 p\ \log^2 q\left(\sum_{\substack{m\le\frac{x}{pq}\\ (m,qp)>1}}\Lambda_{k-1}(mq)\Lambda_{\ell-1}(mp)\right). \qquad (6.14)
\end{aligned}$$

The second term can be estimated using the trivial bound (3.8), exactly as in (6.12), showing that it is $\ll x(\log x)^{k+\ell}$. For the first term, removing the conditions $(m, pq) = 1$ and $q \neq p$ and estimating with the trivial bound (3.8) gives an even smaller error term. This completes the proof of Lemma 6.1.

□

6.2. **Some lemmas.** The following lemmas concerning the Λ_j function and sums over primes are required in the next subsection.

The first Lemma is an improvement on the trivial bound (3.8) for $\Lambda_j(n)$ when n has a known prime factor.

Lemma 6.2. *If $p|m$, then*

$$\Lambda_k(m) \leq k \log p (\log m)^{k-1}.$$

Proof. If $(p, n) = 1$ then

$$\Lambda_k(p^a n) = \sum_{j=1}^{\min(a,k)} \binom{k}{j} \Lambda_j(p^a)\Lambda_{k-j}(n) = \sum_{j=1}^{k} \binom{k}{j}\binom{a-1}{j-1} (\log p)^j \Lambda_{k-j}(n)\,.$$

Setting $a = 1$, we obtain (6.13).

Now

$$\binom{a-1}{j-1} = \frac{(a-1)(a-2)\cdots(a-j+1)}{(j-1)!} \leq \frac{a^{j-1}}{(j-1)!}\,.$$

Hence

$$\begin{aligned}\Lambda_k(p^a n) &\leq \sum_{j=1}^{k} \frac{k}{j}\binom{k-1}{j-1} \frac{a^{j-1}}{(j-1)!} (\log p)^j (\log n)^{k-j} \\ &\leq k \log p \sum_{j=1}^{k} \binom{k-1}{j-1} (\log p^a)^{j-1} (\log n)^{(k-1)-(j-1)}\,.\end{aligned}$$

Setting $j - 1 = i$, this is

$$\begin{aligned}&= k \log p \sum_{i=0}^{k-1} \binom{k-1}{i} (\log p^a)^i (\log n)^{(k-1)-i} \\ &\leq k \log p (\log p^a n)^{k-1}\,.\end{aligned}$$

□

Lemma 6.3. *If $u \geq 2$ and $v \geq 1$ then*

$$\sum_{p \leq x} \frac{\log^u p}{p}\left(\log \frac{x}{p}\right)^v = \frac{(u-1)!v!}{(u+v)!} \log^{u+v} x + O\left(\frac{(u-1)!v!}{(u+v-1)!} \log^{u+v-1} x\right)\,. \tag{6.15}$$

Proof. Let $F(t) = \sum_{p \leq t} \log p/p$, so $F(t) = \log t + E(t)$, where $E(1) = 0$ and $E(t) \ll 1$. We have

$$\begin{aligned}\sum_{p \leq x} \frac{\log^u p}{p}\left(\log \frac{x}{p}\right)^v &= \int_1^x \log^{u-1} t \left(\log \frac{x}{t}\right)^v dF(t) \\ &= \int_1^x \log^{u-1} t \left(\log \frac{x}{t}\right)^v d(\log t) + \int_1^x \log^{u-1} t \left(\log \frac{x}{t}\right)^v dE(t)\end{aligned}$$

$$= I + J,$$

say. In I replace t by x^θ and use Euler's Beta-integral to obtain

$$I = \log^{u+v} x \int_0^1 \theta^{u-1}(1-\theta)^v \, d\theta = \frac{(u-1)!v!}{(u+v)!} \log^{u+v} x, \tag{6.16}$$

which is the main term above. In the error term we integrate by parts and find that

$$J = E(t)\log^{u-1} t \left(\log\frac{x}{t}\right)^v \Big|_1^x - \int_1^x E(t)\left((u-1)\log^{u-2} t\left(\log\frac{x}{t}\right)^v - v\log^{u-1} t\left(\log\frac{x}{t}\right)^{v-1}\right)\frac{dt}{t}. \tag{6.17}$$

The first term vanishes. In the integral make the change of variable $t = x^\theta$ and again use the Beta-integral to obtain

$$\begin{aligned} J &\ll \log^{u+v-1} x \ (u-1)\int_0^1 \theta^{u-2}(1-\theta)^v \, d\theta + \log^{u+v-1} x \ v\int_0^1 \theta^{u-1}(1-\theta)^{v-1}\, d\theta \\ &= 2\frac{(u-1)!v!}{(u+v-1)!}\log^{u+v-1} x, \end{aligned}$$

as claimed. ☐

6.3. **The Estimation of $\sum_{n\le x}\Lambda_k(n)\Lambda_\ell(n)$.** We evaluate the sums over Λ_j which appear in Lemma 6.1

Let

$$\mathcal{S}_{k,l}(x) = \sum_{n\le x}\Lambda_k(n)\Lambda_\ell(n)\,.$$

In this subsection we prove the following theorem.

Theorem 6.4. *If $k > \ell \ge 1$, then*

$$\mathcal{S}_{k,\ell}(x) \ll x\log^{k+\ell-2} x\,. \tag{6.18}$$

If $k \ge 1$, then

$$\mathcal{S}_{k,k}(x) = \frac{k!}{(2k-1)!}x\log^{2k-1} x + O(x\log^{2k-2} x)\,. \tag{6.19}$$

The theorem is proved by induction, using the following proposition.

Proposition 6.5. *For $k \ge \ell \ge 1$ we have*

$$\mathcal{S}_{k,\ell}(x) = \ell\sum_{p\le x}\log^2 p\ \mathcal{S}_{k-1,\ell-1}(x/p) + O(x\log^{k+\ell-2} x). \tag{6.20}$$

Proof. We assume $k \ge \ell$ and begin by unfolding Λ_k in the sum:

$$\begin{aligned} \mathcal{S}_{k,\ell}(x) &= \sum_{n\le x}\left(\sum_{d|n}\Lambda(d)\Lambda_{k-1}\left(\frac{n}{d}\right)\right)\Lambda_\ell(n) \\ &= \sum_{d\le x}\Lambda(d)\left(\sum_{m\le x/d}\Lambda_{k-1}(m)\Lambda_\ell(md)\right) \end{aligned} \tag{6.21}$$

$$
\begin{aligned}
&= \sum_{p^a\le x} \log p \left(\sum_{m\le x/p^a} \Lambda_{k-1}(m)\Lambda_\ell(mp^a) \right)\\
&= \sum_{p\le x} \log p \left(\sum_{m\le x/p} \Lambda_{k-1}(m)\Lambda_\ell(mp) \right)\\
&\qquad + \sum_{\substack{p^a\le x\\ a\ge 2}} \log p \left(\sum_{m\le x/p^a} \Lambda_{k-1}(m)\Lambda_\ell(mp^a) \right)\\
&= \Sigma_1 + \Sigma_2,
\end{aligned}
$$

say. We split Σ_1 into two sums $\Sigma_{1,1}$ and $\Sigma_{1,2}$ according to whether m in the inner sum is or is not coprime to p. By (6.13), if $(m,p)=1$ then $\Lambda_\ell(mp) = \ell\Lambda_{\ell-1}(m)\log p$. Hence

$$
\Sigma_{1,1} = \ell \sum_{p\le x} \log^2 p \left(\sum_{\substack{m\le x/p\\ (m,p)=1}} \Lambda_{k-1}(m)\Lambda_{\ell-1}(m) \right) .
$$

By Lemma 6.2, removing the coprimality condition here introduces a change of

$$
\ll \ell \sum_{p\le x} \log^2 p \left(\frac{x}{p^2} k\ell \log^2 p \log^{k+\ell-4} x \right) \ll x \log^{k+\ell-4} x .
$$

That is,

$$
\Sigma_{1,1} = \ell \sum_{p\le x} \log^2 p \left(\sum_{m\le x/p} \Lambda_{k-1}(m)\Lambda_{\ell-1}(m) \right) + O\left(x \log^{k+\ell-4} x \right) .
$$

For $\Sigma_{1,2}$ we find that

$$
\Sigma_{1,2} = \sum_{p\le x} \log p \left(\sum_{\substack{m\le x/p\\ p|m}} \Lambda_{k-1}(m)\Lambda_l(mp) \right) .
$$

By Lemma 6.2 this is

$$
\begin{aligned}
&\ll \sum_{p\le x} \log p \sum_{r\le x/p^2} \Lambda_{k-1}(rp)\Lambda_\ell(rp^2) \ll k\ell \sum_{p\le x} \log^3 p \sum_{r\le x/p^2} (\log x)^{k+\ell-3}\\
&\ll x(\log x)^{k+\ell-3} \sum_{p\le x} \frac{\log^3 p}{p^2} \ll x(\log x)^{k+\ell-3} .
\end{aligned}
$$

Hence, combining $\Sigma_{1,1}$ and $\Sigma_{1,2}$, we obtain

$$
\Sigma_1 = \ell \sum_{p\le x} \log^2 p \left(\sum_{m\le x/p} \Lambda_{k-1}(m)\Lambda_{\ell-1}(m) \right) + O\left(\ell x (\log x)^{k+\ell-2} \right) , \tag{6.22}
$$

which is the main term in the Proposition.

By the trivial bound (3.8) and Lemma 6.2 we have

$$\begin{aligned}\Sigma_2 &\ll \ell \sum_{\substack{p^a\le x\\ a\ge 2}} \log^2 p \left(\sum_{m\le x/p^a} (\log x)^{k+\ell-2} \right)\\ &\ll \ell x(\log x)^{k+\ell-2} \sum_{\substack{p^a\le x\\ a\ge 2}} \frac{\log^2 p}{p^a}\\ &\ll x(\log x)^{k+\ell-2}\ .\end{aligned}$$

Combining this with (6.22) completes the proof of Proposition 6.5. □

Proof of Theorem 6.4. We first prove the bound (6.18). If $k\ge 2$ then using the basic properties of Λ and Λ_k we have

$$\begin{aligned}\mathcal{S}_{k,1}(x) &= \sum_{n\le x} \Lambda_k(n)\Lambda(n)\\ &= \sum_{p^a\le x} \Lambda_k(p^a)\log p\\ &\le \sum_{\substack{p^a\le x\\ a\ge k}} (\log p^a)^{k+1}\\ &\ll x^{\frac{1}{k}} \log^{k+1} x\\ &\ll x^{\frac{1}{2}+\varepsilon},\end{aligned}$$

which is much smaller than the claimed bound.

Now suppose (6.18) holds for some $\ell>1$ and all $k>\ell$. By Proposition 6.5, the induction hypothesis, and Lemma 6.3:

$$\begin{aligned}\mathcal{S}_{k,\ell+1}(x) &= (\ell+1)\sum_{p\le x} \log^2 p\, \mathcal{S}_{k-1,\ell}\,(x/p) + O(x(\log x)^{k+\ell-1})\\ &\ll \sum_{p\le x} \log^2 p\, \frac{x}{p}\left(\log\frac{x}{p}\right)^{k+\ell-3} + O\left(x(\log x)^{k+\ell-1}\right)\\ &\ll x(\log x)^{k+\ell-1}.\end{aligned}$$

as required. This proves (6.18).

Now we prove (6.19). When $k=1$ we have

$$\mathcal{S}_{1,1}(x) = \sum_{n\le x} \Lambda^2(n) = x\log x + O(x)\ , \tag{6.23}$$

so (6.19) holds in this case. Suppose (6.19) holds for some $k>1$. Then by Proposition 6.5, the induction hypothesis, and Lemma 6.3:

$$\mathcal{S}_{k+1,k+1}(x) = (k+1)\sum_{p\le x} \log^2 p\, \mathcal{S}_{k,k}\left(\frac{x}{p}\right) + O(x\log^{2k} x)$$

$$
\begin{aligned}
&= \frac{(k+1)!}{(2k-1)!} x \sum_{p \le x} \frac{\log^2 p}{p} \left(\log \frac{x}{p} \right)^{2k-1} \tag{6.24}\\
&\quad + O\left(x \sum_{p \le x} \frac{\log^2 p}{p} \left(\log \frac{x}{p} \right)^{2k-2} \right) + O(x \log^{2k} x) \tag{6.25}\\
&= \frac{(k+1)!}{(2k+1)!} x \log^{2k+1} x + O\left(x \log^{2k} x \right), \tag{6.26}
\end{aligned}
$$

as required. □

6.4. **Completion of the proof of Proposition 5.1.** We are now ready to estimate $A_{k,\ell}(x)$.

Proof. Suppose first that $k > \ell \ge 3$. Using Theorem 6.4 and Lemma 6.1 we have

$$
\begin{aligned}
A_{k,\ell}(x) \ll x \sum_{p \le x} \sum_{q \le \frac{x}{p}} \frac{\log^3 p \log^3 q}{pq} \left(\log \frac{x}{pq} \right)^{k+\ell-6}& \\
+ x \sum_{p \le x} \frac{\log^4 p}{p} \left(\log \frac{x}{p} \right)^{k+\ell-4} + O\left(x (\log x)^{k+\ell} \right)& . \tag{6.27}
\end{aligned}
$$

By Lemma 6.3 the first term is

$$
\ll x \sum_{p \le x} \frac{\log^3 p}{p} \left(\log \frac{x}{p} \right)^{k+\ell-3} \ll x \log^{k+\ell} x, \tag{6.28}
$$

and the second term is also $\ll x \log^{k+\ell} x$.

When $k > \ell$ and $\ell = 2$, we obtain the same bound. The only difference is that this time the first term in (6.27) is omitted because $\sum_{n \le x} \Lambda_{k-2}(n) \Lambda_0(n) = \Lambda_{k-2}(1) = 0$ when $k > 2$. This proves (5.7).

Next suppose that $k \ge 3$. By Lemma 6.1 and then Theorem 6.4 we have

$$
\begin{aligned}
A_{k,k}(x) = (k-1)^2 \sum_{p \le x} \sum_{q \le x/p} \log^3 p \log^3 q \sum_{m \le \frac{x}{pq}} \Lambda_{k-2}(m)^2& \\
+ \sum_{p \le x} \log^4 p \sum_{m \le \frac{x}{p}} \Lambda_{k-1}(m)^2 + O\left(x \log^{2k} x \right)& \tag{6.29}\\
= \frac{(k-1)(k-1)!}{(2k-5)!} \sum_{p \le x} \frac{\log^3 p}{p} \sum_{q \le x/p} \frac{\log^3 q}{q} \left(\left(\log \frac{x}{pq} \right)^{2k-5} + O\left(\frac{x}{pq} \log^{2k-6} x \right) \right)& \\
+ \frac{(k-1)!}{(2k-3)!} \sum_{p \le x} \log^4 p \left(\left(\log \frac{x}{pq} \right)^{2k-3} + O\left(\frac{x}{p} \log^{2k-4} x \right) \right) + O\left(x \log^{2k} x \right)& . \tag{6.30}
\end{aligned}
$$

Applying Lemma 6.3 we obtain

$$
A_{k,k}(x) = \frac{2(k-1)(k-1)!}{(2k-2)!} x \sum_{p \le x} \frac{\log^3 p}{p} \left(\log \frac{x}{p} \right)^{2k-2}
$$

$$+\frac{6(k-1)!}{(2k+1)!}x\log^{2k+1}x+O\left(x\log^{2k}x\right)$$

$$=\frac{4(k-1)(k-1)!}{(2k+1)!}x\log^{2k+1}x+\frac{6(k-1)!}{(2k+1)!}x\log^{2k+1}x+O\left(x\log^{2k}x\right)$$

$$=2\frac{(k-1)!}{(2k)!}x\log^{2k+1}x+O\left(x\log^{2k}x\right), \tag{6.31}$$

as claimed.

It remains to do the case $k=2$. The only change in the above analysis is in the first term of (6.29). Since $\Lambda_0(m)=1$ if $m=1$ and 0 otherwise, using Lemma 6.1 and then Theorem 6.4 and Lemma 6.3 we have

$$\begin{aligned}A_{2,2}(x)&=\sum_{p\le x}\log^3 p\sum_{q\le\frac{x}{p}}\log^3 q+\sum_{p\le x}\log^4 p\sum_{m\le x/p}\Lambda(m)^2+O(x\log^4 x)\\&=x\sum_{p\le x}\frac{\log^3 p}{p}\left(\log^2\frac{x}{p}+O\left(\log x\right)\right)+x\sum_{p\le x}\log^4 p\log\frac{x}{p}+O(x\log^4 x)\\&=\frac{2!\,2!}{5!}x\log^5 x+\frac{3!1!}{5!}x\log^5 x+O(x\log^4 x)\\&=\frac{1}{12}x\log^5 x+O(x\log^4 x)\ .\end{aligned}$$

Note that this is the same as the general case with $k=2$. This completes the proof.

Recalling equation (5.2),

$$\mathcal{A}(x)=\sum_{k=0}^{K}A_{k,k}(x)\mathcal{L}^{-2k}+2\sum_{\ell=0}^{K-1}\sum_{\ell<k\le K}A_{k,\ell}(x)\mathcal{L}^{-(k+\ell)}\ , \tag{6.32}$$

and using the fact that $x/\mathcal{L}\ll 1$ if $x=T^\alpha$ with $0<\alpha<C_0$, gives the final statement in Proposition 5.1. □

7. PROOF OF THE EXPLICIT FORMULA

We prove Lemma 3.1 and Proposition 3.2.

7.1. **Proof of Lemma 3.1.**

Proof. Since $\Gamma(s)$ has simple poles at $s=0,-2,-4,\dots$, $L(s)$ has simple poles with residue -1 at $s=-2,-4,\dots$ and a simple pole with residue 1 at $s=1$. It is not difficult to show that $L(s)$ has only simple real zeros at $m_1\approx 7.6, m_2\approx 2.8, m_3\approx -2.6,\dots$, with $m_j\to-\infty$ as $j\to\infty$. Multiplying both sides of (3.1) by $\xi(s)$ and calculating the logarithmic derivative, we obtain

$$\frac{\xi''}{\xi'}(s)=\frac{\xi'}{\xi}(s)+\frac{L'(s)+(\zeta'/\zeta)'(s)}{L(s)+\zeta'/\zeta(s)}\ . \tag{7.1}$$

Suppose now that $\varepsilon>0$. Then there exists an absolute constant C_1 such that for $\sigma\ge 1+\varepsilon$,

$$\left|\frac{\zeta'}{\zeta}(s)\right|\le\frac{\zeta'}{\zeta}(\sigma)\le C_1\varepsilon^{-1} \tag{7.2}$$

and

$$\left|\left(\frac{\zeta'}{\zeta}\right)'(s)\right| \le C_1\varepsilon^{-2}\,. \tag{7.3}$$

Hence, by (3.2), there exists an absolute constant C_2 such that

$$\left|\frac{\zeta'}{\zeta}(s)L(s)^{-1}\right| \le \frac{4C_1}{\varepsilon\log(|s|+2)} < \frac{1}{2} \tag{7.4}$$

for $\sigma \ge 1+\varepsilon$ and $|t| \ge T_\varepsilon = C_2 e^{8C_1/\varepsilon}$. Now let K be an arbitrary large integer. Then by (3.3), (7.1),(7.2), and (7.4),

$$\frac{\xi''}{\xi'}(s) = L(s) + \frac{\zeta'}{\zeta}(s) + \left(\frac{\zeta'}{\zeta}\right)'(s)L(s)^{-1}\sum_{j=0}^{K-1}\left(-\frac{\zeta'}{\zeta}(s)L(s)^{-1}\right)^j + O\left(\frac{1}{\varepsilon^2 2^K}\right) \tag{7.5}$$

for $\sigma \ge 1+\varepsilon$ and $|t| \ge T_\varepsilon = C_2 e^{8C_1/\varepsilon}$. Using the definition of the j-fold von Mangoldt function (3.7), this can be rewritten as

$$\frac{\xi''}{\xi'}(s) = L(s) + \sum_{n=1}^{\infty} n^{-s}\left(-\Lambda(n) + \sum_{j=0}^{K-1}\frac{(\Lambda_j * \Lambda\log)(n)}{L(s)^{j+1}}\right) + O\left(\frac{1}{\varepsilon^2 2^K}\right)\,. \tag{7.6}$$

This completes the proof of Lemma 3.1.

□

7.2. **Comparison to the Hardy Z-function.** We now indicate why the pair correlation functions for the zeros of ξ' and Z' are equal to leading order.

Suppose $\mathcal{X}(s) = \Upsilon(s)\zeta(s)$ and let $\mathcal{L}(s) = \frac{\Upsilon'}{\Upsilon}(s)$. We have

$$\frac{\mathcal{X}''}{\mathcal{X}'}(s) = \mathcal{L}(s) + \frac{\zeta'}{\zeta}(s) + \frac{\mathcal{L}'(s) + (\zeta'/\zeta)'(s)}{\mathcal{L}(s) + \zeta'/\zeta(s)}\,. \tag{7.7}$$

By choosing Υ appropriately, one can obtain either the Riemann ξ-function or the Hardy Z-function. In either case,

$$\mathcal{L}(s) \sim \frac{1}{2}\log s \tag{7.8}$$

and

$$\mathcal{L}'(s) \ll s^{-1}, \tag{7.9}$$

which is all that was used in the calculation of the form factor $F_1(\alpha;T)$.

The lower order terms in $\mathcal{L}(s)$ are different in those two cases, and this should have an effect on the lower order terms of $F_1(\alpha;T)$. Presumably the lower order terms also have an arithmetic component, so both should differ from the analogous expression from random matrix theory.

7.3. **Proof of Proposition 3.2.**

Proof. We will integrate both sides of (3.4) against the following kernel:

$$k(w,s)=\frac{2\sigma-1}{(w-(s-1/2))(w-(1/2-\bar{s}))}\ . \tag{7.10}$$

It is easy to see that

$$k(w,1-\bar{s})=-k(w,s) \tag{7.11}$$

and

$$k(-w,s)=k(w,\bar{s})\ . \tag{7.12}$$

Moreover, as a function of w, $k(w,s)$ has simple poles at $w=s-1/2$ and $w=1/2-\bar{s}$ with residues 1 and -1, respectively. From the partial fraction decomposition for $\zeta'/\zeta(w)$ and the fact that $\zeta(w)$ has $O(\log T)$ zeros with ordinates in the interval $[T,T+1]$, it follows that one can find an increasing, unbounded sequence $\{T_j\}_{j=1}^{\infty}$ such that

$$\frac{\zeta'}{\zeta}(u+\mathrm{i}T_j)\ll\log^2 T_j \tag{7.13}$$

and

$$\left(\frac{\zeta'}{\zeta}\right)'(u+\mathrm{i}T_j)\ll\log^3 T_j \tag{7.14}$$

uniformly for $-1\le u\le 2$. Using these in (3.1) and (7.1), we find that

$$\frac{\xi''}{\xi'}(u+\mathrm{i}T_j)\ll\log^2 T_j \tag{7.15}$$

uniformly for $-1\le u\le 2$. We now write

$$I_j=\frac{1}{2\pi_{\mathrm{i}}}\int_{\mathcal{R}_j}\frac{\xi''}{\xi'}(w+1/2)k(w,s)x^w\mathrm{d}w\ , \tag{7.16}$$

where $x\ge 1, 5/4<\sigma<2$, and $\mathcal{R}_j$ is the positively oriented rectangle with vertices at $c\pm\mathrm{i}T_j, -U\pm\mathrm{i}T_j$, where $c=1/2+\varepsilon$, $\varepsilon<1/8$, and U is a large positive number. The integrand has simple poles at $w=\mathrm{i}\gamma$, $w=s-1/2$, and $w=1/2-\bar{s}$. Now, since $\varepsilon<1/8$, $\mathrm{Re}\,(s-1/2)=\sigma-1/2>3/4>c$. Therefore $s-1/2$ lies outside $\mathcal{R}_j$. Thus, by the calculus of residues,

$$I_j=-\frac{\xi''}{\xi'}(1-\bar{s})x^{1/2-\bar{s}}+\sum_{|\gamma|\le T_j}k(\mathrm{i}\gamma,s)x^{\mathrm{i}\gamma}\ . \tag{7.17}$$

We now estimate the contributions of the horizontal and left edges of $\mathcal{R}_j$ to I_j. Besides (7.15) we need the estimate

$$\frac{\xi''}{\xi'}(w)\ll\log 2|w| \tag{7.18}$$

for $\mathrm{Re}\,w<-1/2$. By (3.1), (3.2), (3.3), and (7.1) this holds in $\mathrm{Re}\,w>3/2$. Hence it holds in *Re* $w<-1/2$ by the functional equation (1.2). By (7.15) and (7.18), the top and bottom edges of $\mathcal{R}_j$ contribute

$$\ll\int_{-U}^{-1}\frac{\log 2|u+iT_j|}{(T_j-t)^2}x^u\mathrm{d}u+\int_{-1}^{c}\frac{\log^2 T_j}{(T_j-t)^2}x^u\mathrm{d}u$$

$$\ll x^{1/2+\varepsilon}\ \frac{\log^2 T_j}{(T_j-t)^2}$$

to I_j. The left edge contributes

$$\begin{aligned}&\ll \int_0^{T_j} \frac{\log\sqrt{U^2+v^2}}{U^2+v^2} x^{-U}\mathrm{d}v\\ &\ll \frac{x^{-U}}{U}\int_0^{\infty}\frac{\log U+\log(1+x^2)}{1+x^2}\,\mathrm{d}x\\ &\ll x^{-U}\frac{\log U}{U}\,.\end{aligned}$$

Letting U and T_j both tend to infinity, we obtain

$$\sum_{\gamma} k(\mathrm{i}\gamma,s)x^{\mathrm{i}\gamma}=\frac{\xi''}{\xi'}(1-\bar{s})x^{1/2-\bar{s}}+\frac{1}{2\pi\mathrm{i}}\int_{c-\mathrm{i}\infty}^{c+\mathrm{i}\infty}\frac{\xi''}{\xi'}(w+\frac{1}{2})k(w,s)x^{w}\mathrm{d}w\,. \tag{7.19}$$

We evaluate the integral here by replacing $\xi''/\xi'(w+1/2)$ by (3.4). Since this representation holds only when $|\operatorname{Im} w|\ge T_\varepsilon$, we have

$$\begin{aligned}&\frac{1}{2\pi\mathrm{i}}\int_{c-\mathrm{i}\infty}^{c+\mathrm{i}\infty}\frac{\xi''}{\xi'}(w+1/2)k(w,s)x^{w}\mathrm{d}w\\ &\qquad=\frac{1}{2\pi\mathrm{i}}\int_{c-\mathrm{i}\infty}^{c+\mathrm{i}\infty}\left(L(w+1/2)+\sum_{n=1}^{\infty}\frac{a_K(n,w+1/2)}{n^{w+1/2}}\right)k(w,s)x^{w}\mathrm{d}w\\ &\qquad\quad+\mathcal{E}_1+\mathcal{E}_2+\mathcal{E}_3\,,\end{aligned} \tag{7.20}$$

where

$$\begin{aligned}\mathcal{E}_1&=\frac{1}{2\pi\mathrm{i}}\int_{c-\mathrm{i}T_\varepsilon}^{c+\mathrm{i}T_\varepsilon}\frac{\xi''}{\xi'}(w+1/2)k(w,s)x^{w}\mathrm{d}w\,,\\ \mathcal{E}_2&=\frac{1}{2\pi\mathrm{i}}\int_{c-\mathrm{i}T_\varepsilon}^{c+\mathrm{i}T_\varepsilon}\left(L(w+1/2)+\sum_{n=1}^{\infty}\frac{a_K(n,w+1/2)}{n^{w+1/2}}\right)k(w,s)x^{w}\mathrm{d}w\,,\\ \mathcal{E}_3&=O\left(\int_{T_\varepsilon}^{\infty}\frac{1}{\varepsilon 2^K}\frac{x^{1/2+\varepsilon}}{1+(v-t)^2}\mathrm{d}v\right)\,.\end{aligned}$$

Now

$$\begin{aligned}\mathcal{E}_1&\ll\log T_\varepsilon\left(\int_0^{T_\varepsilon}\frac{x^{1/2+\varepsilon}}{1+(v-t)^2}\mathrm{d}v\right)\\ &\ll_\varepsilon x^{1/2+\varepsilon}\left(\frac{1}{|t|+2}+\frac{1}{|t-T_\varepsilon|+2}\right)\\ &\ll_\varepsilon \frac{x^{1/2+\varepsilon}}{|t|+2}\,.\end{aligned}$$

Furthermore, by (3.11)

$$\mathcal{E}_2\ll x^{1/2+\varepsilon}\int_0^{T_\varepsilon}\left(\log(v+2)+C_3^{K+2}\sum_{n=1}^{\infty}\frac{(\log n)^{K+2}}{n^{1+\varepsilon}}\right)\frac{dv}{(1+|v-t|)^2}$$

$$\begin{aligned}
&\ll x^{1/2+\varepsilon}\left(\log T_\varepsilon+\left(\frac{C_3}{\varepsilon}\right)^{K+3}\right)\left(\frac{1}{|t|+2}+\frac{1}{|t-T_\varepsilon|+2}\right)\\
&\ll_{\varepsilon,K}\ x^{1/2+\varepsilon}\left(\frac{1}{|t|+2}+\frac{1}{|t-T_\varepsilon|+2}\right)\\
&\ll_{\varepsilon,K}\ \frac{x^{1/2+\varepsilon}}{(|t|+2)}\ .
\end{aligned}$$

Clearly we also have

$$\mathcal{E}_3 \ll_{\varepsilon,K}\ \frac{x^{1/2+\varepsilon}}{(|t|+2)}\ .$$

Therefore,

$$\begin{aligned}
&\frac{1}{2\pi \mathrm{i}}\int_{c-\mathrm{i}\infty}^{c+\mathrm{i}\infty}\frac{\xi''}{\xi'}(w+1/2)k(w,s)x^w\mathrm{d}w\\
&\qquad=\frac{1}{2\pi \mathrm{i}}\int_{c-\mathrm{i}\infty}^{c+\mathrm{i}\infty}\left(L(w+1/2)+\sum_{n=1}^{\infty}\frac{a_K(n,w+1/2)}{n^{w+1/2}}\right)k(w,s)x^w\mathrm{d}w\\
&\qquad\quad+O_{\varepsilon,K}\left(\frac{x^{1/2+\varepsilon}}{|t|+2}\right)\ .
\end{aligned}\tag{7.21}$$

We split the integral on the right-hand side into two parts, namely,

$$\begin{aligned}
\mathcal{I}_1+\mathcal{I}_2=&\frac{1}{2\pi \mathrm{i}}\int_{c-\mathrm{i}\infty}^{c+\mathrm{i}\infty}\left(L(w+1/2)+\sum_{n\le x}\frac{a_K(n,w+1/2)}{n^{w+1/2}}\right)k(w,s)x^w\mathrm{d}w\\
&+\frac{1}{2\pi \mathrm{i}}\int_{c-\mathrm{i}\infty}^{c+\mathrm{i}\infty}\left(\sum_{n>x}\frac{a_K(n,w+1/2)}{n^{w+1/2}}\right)k(w,s)x^w\mathrm{d}w\ .
\end{aligned}\tag{7.22}$$

In $\mathcal{I}_1$ we pull the contour left to $-\infty$ and in doing so we pass a pole of $k(w,s)$ at $w=1/2-\bar{s}$, the poles of $L(w+1/2)$ at $w=1/2,-5/2,-9/2,-13/2,\ldots$, and the poles of $1/L(w+1/2)$ at the points $w=m_3-1/2, m_4-1/2,\ldots$, where the m_j's are the zeros of $L(s)$. We find that

$$\begin{aligned}
\mathcal{I}_1=&-\left(L(-\bar{s})+\sum_{n\le x}\frac{a_K(n,1-\bar{s})}{n^{1-\bar{s}}}\right)x^{1/2-\bar{s}}-\sum_{m=1}^{\infty}k(-2m+1/2,s)x^{-2m+1/2}\\
&+k(1/2,s)x^{1/2}+\sum_{n\le x}\sum_{k=0}^{K}\frac{(\Lambda_k*\Lambda\log)(n)}{\sqrt{n}}\sum_{j=3}^{\infty}\left(\frac{x}{n}\right)^{m_j-1/2}P_{k,j}\left(\log\frac{x}{n},s\right)\ ,
\end{aligned}\tag{7.23}$$

where

$$\begin{aligned}
x^{m_j-1/2}P_{k,j}(\log x,s)&=\mathrm{Res}_{w=m_j-1/2}\,\frac{k(w,s)x^w}{L(w+1/2)^{k+1}}\\
&\ll\frac{(\log x)^k}{k!}x^{m_j-1/2}|k(m_j-1/2,s)|\\
&\ll\frac{(\log x)^k}{k!}x^{m_j-1/2}|m_j-s|^{-2}\ .
\end{aligned}\tag{7.24}$$

Using this and (3.10), we find that the sum over m_j is

$$\begin{aligned}
&\ll \sum_{n\le x} \frac{(\log n)^{K+2}}{\sqrt{n}} \sum_{k=0}^{K} \frac{(\log x/n)^k}{k!} \sum_{j=3}^{\infty} \left(\frac{x}{n}\right)^{m_j-1/2} \frac{1}{|m_j-s|^2} \\
&\ll \sum_{n\le x} \frac{(\log n)^{K+2}}{\sqrt{n}} \sum_{k=0}^{K} \frac{(\log x/n)^k}{k!} \left(\frac{x}{n}\right)^{m_3-1/2} \frac{1}{|s|} \\
&\ll (\log x)^{2K+2} |s|^{-1} x^{m_3-1/2} \sum_{n\le x} \frac{1}{n^{m_3}} \\
&\ll x^{1/2} (\log x)^{2K+2} |s|^{-1} \,.
\end{aligned}$$

The sum over m is

$$\ll \sum_{m=1}^{\infty} \frac{x^{-2m+1/2}}{|2m-s|^2} \ll x^{-3/2}|s|^{-1} \,.$$

Hence

$$\begin{aligned}
\mathcal{I}_1 = &-\left(L(1-\bar{s}) + \sum_{n\le x} \frac{a_K(n,1-\bar{s})}{n^{1-\bar{s}}} \right) x^{1/2-\bar{s}} + k(1/2,s) x^{1/2} \\
&+ O\left(x^{1/2} (\log x)^{2K+2} |s|^{-1} \right) \,.
\end{aligned} \tag{7.25}$$

We treat $\mathcal{I}_2$ by moving the contour away to $+\infty$. This time we pass the pole from $k(w,s)$ at $w = s-1/2$ and the poles of $1/L(w+1/2)$ at $w = m_1 - 1/2$ and $m_2 - 1/2$. Thus we find that

$$\mathcal{I}_2 = -\left(\sum_{n>x} \frac{a_K(n,s)}{n^s} \right) x^{s-1/2} - \sum_{n>x} \sum_{k=0}^{K} \frac{(\Lambda_k * \Lambda \log)(n)}{\sqrt{n}} \sum_{j=1}^{2} \left(\frac{x}{n}\right)^{m_j-1/2} P_{k,j}\left(\log \frac{x}{n}, s \right) \,.$$

We treat the second of these two terms as we did the corresponding term in $\mathcal{I}_1$ and find that it is

$$\begin{aligned}
&\ll \frac{1}{|s|^2} \sum_{n>x} \frac{(\log n)^{K+2}}{\sqrt{n}} \sum_{k=0}^{K} \frac{(\log n/x)^k}{k!} \left(\left(\frac{x}{n}\right)^{m_1-1/2} + \left(\frac{x}{n}\right)^{m_2-1/2} \right) \\
&\ll \frac{1}{|s|^2} \sum_{n>x} \frac{(\log n)^{2K+2}}{\sqrt{n}} \left(\left(\frac{x}{n}\right)^{m_1-1/2} + \left(\frac{x}{n}\right)^{m_2-1/2} \right) \\
&\ll x^{1/2} (\log x)^{2K+2} |s|^{-2} \,.
\end{aligned}$$

Hence,

$$\mathcal{I}_2 = -x^{s-\frac{1}{2}} \left(\sum_{n>x} \frac{a_K(n,s)}{n^s} \right) + O\left(x^{1/2} (\log x)^{2K+2} |s|^{-2} \right) \,. \tag{7.26}$$

Combining (7.19), (7.21), (7.22), (7.25), and (7.26), we see that

$$\sum_{\gamma} k(\mathrm{i}\gamma, s) x^{\mathrm{i}\gamma} = \frac{\xi''}{\xi'}(1-\bar{s}) x^{1/2-\bar{s}} - L(1-\bar{s}) x^{1/2-\bar{s}}$$

$$- x^{-1/2}\left(\sum_{n\le x} a_K(n,1-\bar{s})\left(\frac{x}{n}\right)^{1-\bar{s}} + \sum_{n>x} a_K(n,s)\left(\frac{x}{n}\right)^{s}\right)$$
$$+ k(1/2,s)x^{1/2} + O_{\varepsilon,K}\left(x^{1/2+\varepsilon}|s|^{-1}\right) + O\left(x^{1/2}(\log x)^{2K+2}|s|^{-1}\right) .$$

By the functional equation and (7.1), $\xi''/\xi'(1-\bar{s}) = -\xi''/\xi'(\bar{s}) = -\xi'/\xi(\bar{s}) + O(1)$ for $5/4 < \sigma < 2$. By (3.2) and (3.1) this equals $-1/2\log\bar{s} + O(1)$. Hence, writing $\tau = |t| + 2$, and using the definition (7.10), we finally obtain the explicit formula in Proposition 3.2. ☐

REFERENCES

[1] *M. Berry*, Universal oscillations of high derivatives. Proc. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci. 461 (2005), no. 2058, 1735–1751.

[2] *B. Conrey*, More that two-fifths of the zeros of the Riemann zeta-function are on the critical line. J. Reine Angew. Math. 399 (1989), 1–26

[3] *B. Conrey*, *D. Farmer*, *J. Keating*, *M. Rubinstein*, and *N. Snaith*, Integral moments of L-functions. Proc. London Math. Soc. (3) 91 (2005), no. 1, 33–104.

[4] *B. Conrey* and *H. Iwaniec*, Spacing of zeros of Hecke L-functions and the class number problem. Acta Arith. 103 (2002), no. 3, 259–312.

[5] *B. Conrey* and *Soundararajan*, personal communication.

[6] *T. Craven*, *G. Csordas*, and *W. Smith*. The zeros of derivatives of entire functions and the Pólya-Wiman conjecture. Ann. of Math. (2) 125 (1987), no. 2, 405431.

[7] *D. Farmer*, Counting distinct zeros of the Riemann zeta-function. Electron. J. Combin. 2 (1995), Research Paper 1, approx. 5 pp.

[8] *D. Farmer* and *R. Rhoades*, Differentiation evens out zero spacings. Trans. Amer. Math. Soc. 357 (2005), no. 9, 3789–3811

[9] *D. Farmer* and *S. Gonek*, Mean values of ζ'/ζ, the distribution of almost primes, and the distribution of zeros. Preprint.

[10] *D. Farmer* and *M. Yerrington*, Crystallization of random trigonometric polynomials. J. Stat. Phys. 123 (2006), no. 6, 1219–1230

[11] *D. A. Goldston* and *H. L. Montgomery*, Pair correlation of zeros and primes in short intervals, Analytic Number Theory and Diophantine Problems, Birkhauser, Boston, Mass. (1987), 183–203.

[12] *D. Hejhal*, On the triple correlation of zeros of the zeta function. Internat. Math. Res. Notices 1994, no. 7, 293ff.,

[13] *H. Ki*, The Riemann Ξ-function under repeated differentiation. J. Number Theory 120 (2006), no. 1, 120–131.

[14] *H. Ki* and *Y.-O. Kim*, On the number of nonreal zeros of real entire functions and the Fourier-Pólya conjecture. Duke Math. J. 104 (2000), no. 1, 4573.

[15] *Y.-O. Kim*, A proof of the Pólya-Wiman conjecture. Proc. Amer. Math. Soc. 109 (1990), no. 4, 10451052.

[16] *H.L. Montgomery*, The pair correlation of zeros of the zeta function, Analytic Number Theory, Proceedings of Symposia in Pure Mathematics **2**4 (1973) 181–193

[17] *H.L. Montgomery* and *K. Soundararajan*, Beyond pair correlation. Paul Erds and his mathematics, I (Budapest, 1999), 507–514, Bolyai Soc. Math. Stud., 11, Jnos Bolyai Math. Soc., Budapest, 2002.

[18] *Z. Rudnick* and *P. Sarnak*, Zeros of principal L-functions and random matrix theory. Duke Math. J. 81 (1996), no. 2, 269–322.

[19] E.C. Titchmarsh, *The Theory of the Riemann Zeta-Function* (2nd edition, revised by D.R. Heath-Brown), Oxford Science Publications, 1986

AMERICAN INSTITUTE OF MATHEMATICS, 360 PORTAGE AVENUE, PALO ALTO, CA 94306-2244, USA.
E-mail address: `farmer@aimath.org`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ROCHESTER, ROCHESTER, NY 14627, USA
E-mail address: `gonek@math.rochester.edu`